

Riemann–Hilbert analysis for Laguerre polynomials with large negative parameter

A.B.J. Kuijlaars¹ and K.T-R McLaughlin²

Abstract. We study the asymptotic behavior of Laguerre polynomials $L_n^{(\alpha_n)}(nz)$ as $n \rightarrow \infty$, where α_n is a sequence of negative parameters such that $-\alpha_n/n$ tends to a limit $A > 1$ as $n \rightarrow \infty$. These polynomials satisfy a non-hermitian orthogonality on certain contours in the complex plane. This fact allows the formulation of a Riemann–Hilbert problem whose solution is given in terms of these Laguerre polynomials. The asymptotic analysis of the Riemann–Hilbert problem is carried out by the steepest descent method of Deift and Zhou, in the same spirit as done by Deift et al. for the case of orthogonal polynomials on the real line. A main feature of the present paper is the choice of the correct contour.

Keywords: Riemann–Hilbert problems, generalized Laguerre polynomials, strong asymptotics, steepest descent method.

2000 Mathematics Subject Classification: 30E15, 33C45.

1 Introduction

1.1 Generalized Laguerre polynomials

The classical Laguerre polynomials $L_n^{(\alpha)}$ are orthogonal on the interval $[0, \infty)$ with respect to the weight $x^\alpha e^{-x}$, that is

$$\int_0^\infty L_n^{(\alpha)}(x) L_m^{(\alpha)}(x) x^\alpha e^{-x} dt = 0, \quad \text{if } n \neq m. \quad (1.1)$$

The integral in (1.1) converges only if $\alpha > -1$. The Laguerre polynomials are given by the explicit formula

$$L_n^{(\alpha)}(x) = \sum_{k=0}^n \binom{n+\alpha}{n-k} \frac{(-x)^k}{k!} \quad (1.2)$$

and by the Rodrigues formula

$$L_n^{(\alpha)}(x) = \frac{1}{n!} x^{-\alpha} e^x \left(\frac{d}{dx} \right)^n [x^{\alpha+n} e^{-x}], \quad (1.3)$$

see e.g. [30]. Both (1.2) and (1.3) make sense for arbitrary α (even complex) and they define the generalized non-classical Laguerre polynomials. The recurrence relation

$$-x L_n^{(\alpha)}(x) = (n+1) L_{n+1}^{(\alpha)}(x) - (2n+\alpha+1) L_n^{(\alpha)}(x) + (n+\alpha) L_{n-1}^{(\alpha)}(x), \quad (1.4)$$

with $L_0^{(\alpha)} \equiv 1$, $L_{-1}^{(\alpha)} \equiv 0$, and the second order differential equation

$$xy''(x) + (\alpha+1-x)y'(x) + ny(x) = 0, \quad y(x) = L_n^{(\alpha)}(x), \quad (1.5)$$

continue to hold for arbitrary α . We consider in this paper only real and negative α , although extensions to complex α are possible.

¹Supported in part by FWO research project G.0176.02, by INTAS project 00-272, and by a research grant of the Fund for Scientific Research–Flanders.

²Supported in part by NSF grant # DMS-9970328.

1.2 Relations to other polynomials

Laguerre polynomials with negative parameters appear in the literature in a number of forms. First we note the special cases $\alpha = -n$ and $\alpha = -n - 1$, where we have

$$L_n^{(-n)}(z) = \frac{(-1)^n}{n!} z^n, \quad (1.6)$$

and

$$L_n^{(-n-1)}(z) = (-1)^n \sum_{k=0}^n \frac{z^k}{k!}, \quad (1.7)$$

respectively. Thus for $\alpha = -n - 1$, the generalized Laguerre polynomial agrees (up to a sign if n is odd) with the partial sum of the exponential series. More generally, we have that for $\alpha = -n - m$ with $n, m \in \mathbb{N}$, the generalized Laguerre polynomials appear as the numerator and denominator polynomials in the rational Padé approximant for the exponential function. To be precise, if

$$p_{n,m}(x) = (-1)^n \binom{n+m}{n}^{-1} L_n^{(-n-m-1)}(x) \quad (1.8)$$

and

$$q_{n,m}(x) = (-1)^m \binom{n+m}{m}^{-1} L_m^{(-n-m-1)}(x) \quad (1.9)$$

then $p_{n,m}(0) = q_{n,m}(0) = 1$, and

$$p_{n,m}(x) - q_{n,m}(x)e^x = O(x^{n+m+1}) \quad \text{as } x \rightarrow 0, \quad (1.10)$$

see e.g. [23], [25].

Laguerre polynomials with negative parameters are also related to the so-called generalized Bessel polynomials

$$y_n(z; a) = \sum_{k=0}^n \binom{n}{k} (n+a-1)_k \left(\frac{z}{2}\right)^k, \quad (1.11)$$

since

$$y_n(z; a) = (-1)^n n! \left(\frac{z}{2}\right)^n L_n^{(-2n-a+1)}\left(\frac{2}{z}\right). \quad (1.12)$$

The usual Bessel polynomials correspond to $a = 2$ in (1.11). See Grosswald [17] for a comprehensive account of these polynomials.

1.3 Earlier work on asymptotics

For $\alpha > -1$, the Laguerre polynomials satisfy the orthogonality (1.1) on the positive real axis, and therefore they have only positive real zeros. This property is lost for $\alpha < -1$. Indeed, the generalized Laguerre polynomials may have many non-real zeros. In Figure 1 we have plotted the zeros of $L_{40}^{(-40A)}(40z)$ for a number of values of $A > 0$. Similar plots are shown in the paper [20].




Figure 1: Zeros of generalized Laguerre polynomials $L_n^{(-An)}(nx)$ for $n = 40$ and $A = 0.81$ (left), $A = 1.01$ (middle), and $A = 2$ (right).

In Figure 1 we see that the zeros cluster along certain curves in the complex plane. Martínez et al. [20] identified these curves as trajectories of a quadratic differential, depending on a parameter A . For $A > 1$, the curve is a simple arc, which as A decreases to 1 closes itself to form for $A = 1$ the well-known Szegő curve [29], [24]. For $0 < A < 1$, the curve consists of a closed loop together with an interval on the positive real axis.

A number of rigorous results on the asymptotic behavior of the zeros of generalized Laguerre polynomials $L_n^{(\alpha_n)}(nx)$ such that

$$\lim_{n \rightarrow \infty} \frac{\alpha_n}{n} = -A \quad (1.13)$$

are known from the literature. The first result is due to Szegő [29] who studied the partial sum of the exponential series, that is $\alpha_n = -n - 1$ see (1.7). Szegő showed that the normalized zeros tend to the curve which now bears his name. Olver [21] considered the zeros of Hankel functions, which includes the Bessel polynomials as a special case. In terms of the generalized Laguerre polynomials, this is the case $\alpha_n = -2n - 1$. Saff and Varga [26] studied the zeros and poles of Padé approximants to the exponential function, see (1.8)–(1.9). Their main result says that for integers $\alpha_n < -n$ such that (1.13) holds, all zeros of $L_n^{(\alpha_n)}(nx)$ tend to a well-defined curve as $n \rightarrow \infty$. The curve depends on $A \geq 1$ only, and coincides with the curve described in [20]. Saff and Varga also obtained the weak limit of the zero counting measures. The proofs in [26] can be extended without any difficulty to non-integer $\alpha_n < -n$.

Stated in terms of generalized Bessel polynomials (1.11)–(1.12) asymptotic results on zeros are due to De Bruin et al. [4], Carpenter [5], and Wong and Zhang [31]. These papers deal with the limit (1.13) with $A = 2$. The latter paper also presents uniform asymptotic expansions of the generalized Bessel polynomials. In very recent work, Dunster [14] establishes uniform asymptotic expansions in the complex plane for the case of general A , with the exception of $A = 0$ and $A = 1$. The results on zeros in [14] are restricted to the case $A > 1$.

1.4 Asymptotics from Riemann–Hilbert problems

Most papers cited above use some form of the steepest descent technique for integrals, see especially [26] and [31]. Martínez et al. [20] use an orthogonality relation in the complex plane satisfied by generalized Laguerre polynomials. The approach of Dunster [14] starts from the differential equation (1.5) and is based on techniques developed by Olver [22].

In this paper we derive asymptotics of generalized Laguerre polynomials using the non-linear steepest descent / stationary phase method for Riemann–Hilbert problems introduced by Deift and Zhou in [12], and further developed in [13] and [11]. In later developments,

the method was applied successfully to problems in random matrix theory [8], [9], and in orthogonal polynomials [3], [9], [10], [19], and combinatorics [2]. For review of some of these developments, and a pedagogic introduction to some of the material of random matrix theory, orthogonal polynomials, and Riemann–Hilbert problems, see [6].

The Riemann–Hilbert approach to the asymptotics of generalized Laguerre polynomials starts from the observation that these polynomials satisfy orthogonality relations in the complex plane. The orthogonality is on a contour Σ going around the positive real axis, but otherwise being quite arbitrary. The orthogonality property allows the formulation of a Riemann–Hilbert problem, due to Fokas, Its, and Kitaev [15], whose solution is given in terms of $L_n^{(\alpha)}$. The Riemann–Hilbert problem is analyzed in the large n limit with the steepest descent method as done in [9] and [10] for orthogonal polynomials on the real line.

A novel feature for the problem at hand is that the arbitrary contour Σ has to be chosen in a correct way in order to arrive at a Riemann–Hilbert problem which is amenable to subsequent asymptotic analysis. The correct contour was described in [20]. It is a curve with the S-property of Stahl [27] and Gonchar and Rakhmanov [16]. The structure of the curve depends on the value of A as already explained before. In this paper we analyze the case of an open contour, that is, the case $A > 1$. In subsequent work we consider the case of a closed loop plus an interval ($0 < A < 1$) and the case of a single closed contour ($A = 1$). We note that the steepest descent / stationary phase method for Riemann–Hilbert problems was augmented to handle cases in which the contour selection involves determining a set of nontrivial curves in the plane in [18], by Kamvissis, McLaughlin, and Miller, in the context of the semi-classical limit of the focusing nonlinear Schrödinger equation.

We emphasize that the main interest in the present paper lies in the method we use and not in the results obtained for the Laguerre polynomials. In particular, we do not improve upon the asymptotic expansions of Dunster [14]. The steepest descent method for Riemann–Hilbert problems is a very powerful new method, and its use in the study of classical special functions is new. In future work we consider generalized Laguerre polynomials for the cases $0 < A < 1$ and $A = 1$, and the Riemann–Hilbert approach will lead to new results for these cases.

2 Complex orthogonality and the formulation of the Riemann–Hilbert problem

2.1 Orthogonality

For $\alpha < -1$, the generalized Laguerre polynomial $L_n^{(\alpha)}$ is not orthogonal on the positive real axis, but instead satisfies a non-hermitian orthogonality in the complex plane.

Let $\mathcal{F}$ be the collection of all simple Jordan curves Σ in $\mathbb{C} \setminus [0, \infty)$ that are symmetric with respect to the real axis, and such that there is $M > 0$ such that for all $x \geq M$, there is $y(x) > 0$, such that the intersection of Σ with $\operatorname{Re} z = x$ consists of the two points $x \pm iy(x)$, and $\lim_{x \rightarrow \infty} y(x) = L$ exists and is finite (possibly 0). Any curve $\Sigma \in \mathcal{F}$ divides the complex plane into two domains, Ω_+ and Ω_- , where Ω_- contains the positive real axis. We choose the orientation of Σ such that Ω_+ is on the $+$ -side (i.e., on the left) while traversing Σ and Ω_- is on the $-$ -side. So Σ is oriented clockwise as in Figure 2.

In what follows we define x^α with a branch cut along the positive real axis. Thus $x^\alpha = |x|^\alpha e^{i\alpha \arg x}$ with $\arg x \in [0, 2\pi)$.




Figure 2: Example of contour Σ

Lemma 2.1 *Let $\Sigma \in \mathcal{F}$, $n \in \mathbb{N}$, and $\alpha \in \mathbb{R}$. Then*

$$\int_{\Sigma} L_n^{(\alpha)}(x) x^k x^{\alpha} e^{-x} dx = 0, \quad \text{for } k = 0, 1, \dots, n-1. \quad (2.1)$$

If in addition $\alpha + n + 1 \notin \mathbb{N}$, then

$$\int_{\Sigma} L_n^{(\alpha)}(x) x^k x^{\alpha} e^{-x} dx \neq 0, \quad \text{for } k = n. \quad (2.2)$$

Proof. The orthogonality (2.1) follows from the Rodrigues formula (1.3) by repeated integration by parts. In the same way, we also get

$$\int_{\Sigma} L_n^{(\alpha)}(x) x^n x^{\alpha} e^{-x} dx = \frac{1}{n!} \int_{\Sigma} \left(\frac{d}{dx} \right)^n (x^{\alpha+n} e^{-x}) x^n dx = (-1)^n \int_{\Sigma} x^{\alpha+n} e^{-x} dx$$

For $\alpha + n > -1$, we deform Σ to the positive real axis to obtain

$$\begin{aligned} \int_{\Sigma} L_n^{(\alpha)}(x) x^n x^{\alpha} e^{-x} dx &= (-1)^n (1 - e^{2\pi i \alpha}) \int_0^{\infty} x^{\alpha+n} e^{-x} dx \\ &= (-1)^{n+1} 2i e^{\pi i \alpha} \sin(\pi \alpha) \Gamma(\alpha + n + 1), \end{aligned} \quad (2.3)$$

where Γ denotes the Gamma function. By analytic continuation the integral in (2.2) is equal to (2.3) for every α , and (2.2) follows. $\square$

The formula (2.1) expresses orthogonality with respect to the complex measure $x^{\alpha} e^{-x} dx$ on Σ .

2.2 Riemann–Hilbert problem

Let $\alpha \in \mathbb{R}$. We consider the monic polynomials

$$P_n(z) = \frac{(-1)^n n!}{n^n} L_n^{(\alpha)}(nz), \quad n = 0, 1, \dots, \quad (2.4)$$

Introducing a change of variables $x = nz$ in (2.1) and (2.2), we see that

$$\int_{\Sigma} P_n(z) z^k z^{\alpha} e^{-nz} dz \begin{cases} = 0, & \text{for } k = 0, 1, \dots, n-1, \\ \neq 0, & \text{for } k = n, \end{cases} \quad (2.5)$$

for every contour $\Sigma \in \mathcal{F}$, provided that $\alpha + n + 1 \notin \mathbb{N}$.

The polynomial P_n is characterized through a Riemann–Hilbert problem due to Fokas, Its, and Kitaev [15].

Riemann–Hilbert problem for Y :

Let Σ be a contour from the class $\mathcal{F}$, that divides the complex plane into two parts Ω_+ and Ω_- , as above. The problem is to determine a 2×2 matrix valued function $Y : \mathbb{C} \setminus \Sigma \rightarrow \mathbb{C}^{2 \times 2}$ such that the following hold.

- (a) $Y(z)$ is analytic for $z \in \mathbb{C} \setminus \Sigma$,
- (b) $Y(z)$ possesses continuous boundary values for $z \in \Sigma$, denoted by $Y_+(z)$ and $Y_-(z)$, where $Y_+(z)$ and $Y_-(z)$ denote the limiting values of $Y(z')$ as z' approaches $z \in \Sigma$ from Ω_+ and Ω_- , respectively, and

$$Y_+(z) = Y_-(z) \begin{pmatrix} 1 & z^{\alpha} e^{-nz} \\ 0 & 1 \end{pmatrix}, \quad \text{for } z \in \Sigma, \quad (2.6)$$

- (c) $Y(z)$ has the following behavior as $z \rightarrow \infty$:

$$Y(z) = \left(I + O\left(\frac{1}{z}\right) \right) \begin{pmatrix} z^n & 0 \\ 0 & z^{-n} \end{pmatrix} \quad \text{as } z \rightarrow \infty, \quad z \in \mathbb{C} \setminus \Sigma. \quad (2.7)$$

Proposition 2.2 *Let $\alpha \in \mathbb{R}$ with $\alpha + n \notin \mathbb{N}$. Then the unique solution of the Riemann–Hilbert problem for Y is given by*

$$Y(z) = \begin{pmatrix} P_n(z) & \frac{1}{2\pi i} \int_{\Sigma} \frac{P_n(\zeta) \zeta^{\alpha} e^{-n\zeta}}{\zeta - z} d\zeta \\ Q_{n-1}(z) & \frac{1}{2\pi i} \int_{\Sigma} \frac{Q_{n-1}(\zeta) \zeta^{\alpha} e^{-n\zeta}}{\zeta - z} d\zeta \end{pmatrix} \quad (2.8)$$

where $P_n(z)$ is the monic generalized Laguerre polynomial (2.4) and

$$Q_{n-1}(z) = \frac{(-1)^{n+1} n^{n+\alpha} \pi e^{-\pi i \alpha}}{\sin(\pi \alpha) \Gamma(\alpha + n)} L_{n-1}^{(\alpha)}(nz). \quad (2.9)$$

Proof. The proof is as in [6, Section 3.2]. Here we will only give the proof for the second row, since that is where the condition on α plays a role.

From (2.6) it follows that Y_{21} is an entire function, which by (2.7) satisfies $Y_{21}(z) = O(z^{n-1})$ as $z \rightarrow \infty$. Therefore $Y_{21}(z) = Q_{n-1}(z)$ for some polynomial Q_{n-1} of degree at most $n-1$. The (2,2) entry of the condition (2.6) is $(Y_{22})_+(z) = (Y_{22})_-(z) + Q_{n-1}(z) z^{\alpha} e^{-nz}$, which by the Sokhotskii–Plemelj formula yields

$$Y_{22}(z) = \frac{1}{2\pi i} \int_{\Sigma} \frac{Q_{n-1}(\zeta) \zeta^{\alpha} e^{-n\zeta}}{\zeta - z} d\zeta. \quad (2.10)$$

From the (2,2) entry of (2.7) it follows that $Y_{22}(z) = z^{-n} + O(z^{-n-1})$ as $z \rightarrow \infty$. Because of (2.10) this gives the conditions

$$\int_{\Sigma} Q_{n-1}(\zeta) \zeta^{\alpha} e^{-n\zeta} \zeta^k d\zeta = 0 \quad \text{for } k = 0, \dots, n-2, \quad (2.11)$$

and

$$\int_{\Sigma} Q_{n-1}(\zeta) \zeta^{\alpha} e^{-n\zeta} \zeta^{n-1} d\zeta = -2\pi i. \quad (2.12)$$

The orthogonality conditions (2.11) are satisfied if $Q_{n-1}(z) = d_n L_{n-1}^{(\alpha)}(nz)$, and the constant d_n must be chosen so that (2.12) holds as well. Because of Lemma 2.1 this can be done if $\alpha + n \notin \mathbb{N}$, and the result is given by (2.9). $\square$

3 Selection of the right contour

3.1 The right contour Σ

In Section 3–6, we assume $n \in \mathbb{N}$ and $\alpha < -n$ are fixed. We write $A = -\alpha/n$, so that $A > 1$. [Later, when we let $n \rightarrow \infty$, α and A , as well as other notions introduced in these sections, will depend on n .]

A major step in the analysis of the Riemann–Hilbert problem for Y is the selection of the right contour. In order that the subsequent analysis works, the contour cannot be arbitrary but has to be chosen in a precise way. The contour depends on A . We define

$$\beta = 2 - A + 2i\sqrt{A-1} \quad (3.1)$$

Martinez et al. [20] showed that the values of z for which

$$\frac{1}{2\pi i} \int_{\bar{\beta}}^z \frac{(s-\beta)^{1/2}(s-\bar{\beta})^{1/2}}{s} ds \quad \text{is real,}$$

where the branch of the square roots is chosen so that they are analytic and single valued on the path of integration from $\bar{\beta}$ to z , form a system of curves as shown in Figure 3 for a number of values of A . In geometric function theory these curves are known as trajectories of the quadratic differential $\frac{(s-\beta)(s-\bar{\beta})}{s^2} d^2s$, see [28]. We see four smooth (in fact analytic)




Figure 3: Curves where $\frac{1}{2\pi i} \int_{\bar{\beta}}^z \frac{R(s)}{s} ds$ is real, for the values $A = 1.01$ (left) and $A = 2$ (right). The two points of intersection are $\bar{\beta}$ and β .

curves. Two curves are connecting $\bar{\beta}$ with β , one of them crosses the negative real axis, and the other one crosses the positive real axis.

Definition 3.1 We define Γ as the trajectory of the quadratic differential $\frac{(s-\beta)(s-\bar{\beta})}{s^2} d^2s$ from $\bar{\beta}$ to β which crosses the negative real axis. Γ is oriented from $\bar{\beta}$ to β .

We put

$$R(z) = (z - \beta)^{1/2}(z - \bar{\beta})^{1/2}, \quad z \in \mathbb{C} \setminus \Gamma, \quad (3.2)$$

where the branch is chosen which is defined and analytic on $\mathbb{C} \setminus \Gamma$, and which is such that $R(z) \sim z$ as $z \rightarrow \infty$. For $s \in \Gamma$, we use $R_+(s)$ and $R_-(s)$ to denote the limits from the $+$ -sides and $-$ -sides, respectively. As usual, the $+$ -side of an oriented curve lies to the left, and the $-$ -side lies to the right, if one traverses the curve.

Then by definition of Γ , we have

$$\frac{1}{2\pi i} \int_{\bar{\beta}}^z \frac{R_+(s)}{s} ds \quad \text{is real for every } z \in \Gamma, \quad (3.3)$$

with integration along the $+$ -side of Γ .

It is also of interest to know where $\frac{1}{2\pi i} \int_{\bar{\beta}}^z \frac{R(s)}{s} ds$ or $\frac{1}{2\pi i} \int_{\beta}^z \frac{R(s)}{s} ds$ is purely imaginary. These are the dotted curves shown in Figure 4. The dotted curves are analytic extensions of the solid ones.




Figure 4: Curves where $\frac{1}{2\pi i} \int_{\bar{\beta}}^z \frac{R(s)}{s} ds$ is real (solid lines) and curves where $\frac{1}{2\pi i} \int_{\bar{\beta}}^z \frac{R(s)}{s} ds$ or $\frac{1}{2\pi i} \int_{\beta}^z \frac{R(s)}{s} ds$ is purely imaginary (dotted lines), for the values $A = 1.01$ (left) and $A = 2$ (right).

We can now state which contour Σ to choose.

Definition 3.2 We let Σ be the contour in $\mathcal{F}$ consisting of Γ together with the two dotted curves that form the analytic extension of Γ .

We denote the part of $\Sigma \setminus \Gamma$ in the lower half-plane by Σ_1 and its mirror image in the upper half-plane by Σ_2 .

So we have a disjoint union $\Sigma = \Gamma \cup \Sigma_1 \cup \Sigma_2$. Figure 5 shows the curve Σ for two values of A , together with the zeros of the corresponding Laguerre polynomial $L_{40}^{(-40A)}(40x)$ of degree 40. The figure shows that the zeros are close to Γ , and that they are in the domain Ω_- . These findings will be confirmed by our final result, Corollary 7.2 below.

Remark 3.3 We have chosen Σ_2 so that $\int_{\beta}^z \frac{R(s)}{s} ds$ is real and positive on Σ_2 . This is not essential. What is important for the subsequent analysis is that it has positive real part on Σ_2 . This means that we have the freedom to deform Σ_2 , as long as we take care that the real part of $\int_{\beta}^z \frac{R(s)}{s} ds$ is positive on Σ_2 . However, it will be convenient to do this deformation only away from β , so that in a neighborhood of β , we have Σ_2 exactly as we defined it.

Similar remarks apply to $\int_{\bar{\beta}}^z \frac{R(s)}{s} ds$ and Σ_1 .




Figure 5: The curves $\Sigma = \Gamma \cup \Sigma_1 \cup \Sigma_2$, and the zeros of $L_{40}^{(-40A)}(40x)$, for the values $A = 1.01$ (left) and $A = 2$ (right).

3.2 A probability measure on Γ

The following proposition gives one of the crucial properties of Γ .

Proposition 3.4 *The (complex) measure $\frac{1}{2\pi i} \frac{R_+(s)}{s} ds$ is a probability measure on Γ .*

Proof. We show first that

$$\frac{1}{2\pi i} \int_{\Gamma} \frac{R_+(s)}{s} ds = 1. \quad (3.4)$$

Let $I = \int_{\Gamma} \frac{R_+(s)}{s} ds$. Since $R_-(s) = -R_+(s)$ for $s \in \Gamma$, we have

$$2I = \int_{\Gamma} \left(\frac{R_+(s)}{s} - \frac{R_-(s)}{s} \right) ds = \oint_{\gamma} \frac{R(s)}{s} ds,$$

where γ is a closed contour encircling the curve Γ once in the clockwise direction and not encircling $z = 0$.

After contour deformation, we pick up residues at $z = 0$ and at $z = \infty$, namely

$$2I = 2\pi i \operatorname{Res}_{z=0} \frac{R(z)}{z} - 2\pi i \operatorname{Res}_{z=\infty} \frac{R(z)}{z}. \quad (3.5)$$

The residue at $z = 0$ is

$$\operatorname{Res}_{z=0} \frac{R(z)}{z} = R(0) = |\beta| = \sqrt{(2-A)^2 + 4(A-1)} = A \quad (3.6)$$

and the residue at $z = \infty$ is the coefficient of z^{-1} in the Laurent expansion of $R(z)/z$

$$\begin{aligned} \frac{R(z)}{z} &= (1 - \beta/z)^{1/2} (1 - \bar{\beta}/z)^{1/2} \\ &= \left(1 - \frac{\beta}{2z} + O(z^{-2}) \right) \left(1 - \frac{\bar{\beta}}{2z} + O(z^{-2}) \right) \\ &= 1 - \frac{\beta + \bar{\beta}}{2z} + O(z^{-2}). \end{aligned}$$

Thus

$$\operatorname{Res}_{z=\infty} \frac{R(z)}{z} = -\frac{\beta + \bar{\beta}}{2} = -\operatorname{Re} \beta = -(2-A). \quad (3.7)$$

Hence by (3.5)–(3.7) we have $2I = 2\pi i(A + (2 - A)) = 4\pi i$, so that (3.4) follows.

Having (3.4) we can now prove the proposition. Let $t \mapsto z(t)$ for $t \in [0, t_0]$, be the arc length parametrization of Γ starting at $\bar{\beta}$. Thus $z(0) = \bar{\beta}$ and $z(t_0) = \beta$. Then we have for $z = z(t)$ with $t \in (0, t_0)$,

$$\frac{1}{2\pi i} \int_{\bar{\beta}}^z \frac{R_+(s)}{s} ds = \frac{1}{2\pi i} \int_0^t \frac{R_+(z(\tau))}{z(\tau)} z'(\tau) d\tau, \quad (3.8)$$

and this is real for every t by construction of Γ . It has the value 0 for $t = 0$ and the value 1 for $t = t_0$ (due to (3.4)). The derivative of (3.8) with respect to t is $\frac{1}{2\pi i} \frac{R_+(z(t))}{z(t)} z'(t)$ which is not zero for $t \in (0, t_0)$. Thus (3.8) can only increase from 0 to 1 as t increases from 0 to t_0 . Hence $\frac{1}{2\pi i} \frac{R_+(s)}{s} ds$ is a positive measure on Γ . It is a probability measure because of (3.4). $\square$

3.3 Auxiliary functions

With the measure $d\mu(s) = \frac{1}{2\pi i} \frac{R_+(s)}{s} ds$ on Γ we define the so-called g -function as follows.

Definition 3.5 The g -function is the complex logarithmic potential of μ , that is,

$$g(z) = \int_{\Gamma} \log(z - s) d\mu(s), \quad z \in \mathbb{C} \setminus (\Gamma \cup \Sigma_1), \quad (3.9)$$

where for each s we view $\log(z - s)$ as an analytic function of the variable z , with branch cut emanating from $z = s$. The cut is taken along $\Gamma \cup \Sigma_1$.

We need two more functions.

Definition 3.6 The ϕ -function is defined as

$$\phi(z) = \frac{1}{2} \int_{\beta}^z \frac{R(s)}{s} ds, \quad z \in \mathbb{C} \setminus (\Gamma \cup \Sigma_1 \cup [0, \infty)), \quad (3.10)$$

where the path of integration from β to z lies entirely in $\mathbb{C} \setminus (\Gamma \cup \Sigma_1 \cup [0, \infty))$, except for the initial point β .

The $\tilde{\phi}$ -function is defined as

$$\tilde{\phi}(z) = \frac{1}{2} \int_{\bar{\beta}}^z \frac{R(s)}{s} ds, \quad z \in \mathbb{C} \setminus (\Gamma \cup \Sigma_2 \cup [0, \infty)), \quad (3.11)$$

where the path of integration from $\bar{\beta}$ to z lies entirely in $\mathbb{C} \setminus (\Gamma \cup \Sigma_2 \cup [0, \infty))$, except for the initial point $\bar{\beta}$.

It is immediate from (3.3) that $\tilde{\phi}_+(z)$ is purely imaginary for $z \in \Gamma$. By Proposition 3.4 its imaginary part increases from 0 to π as z traverses the curve Γ from $\bar{\beta}$ to β . In particular we have $\tilde{\phi}_+(\beta) = \pi i$. Similarly $\tilde{\phi}_-(\beta) = -\pi i$. Thus we have

$$\tilde{\phi}(z) = \begin{cases} \phi(z) + \pi i & \text{for } z \in \Omega_+, \\ \phi(z) - \pi i & \text{for } z \in \Omega_-. \end{cases} \quad (3.12)$$

Proposition 3.7 *There is a constant ℓ such that*

$$g(z) = \frac{1}{2} (A \log z + z + \ell) - \phi(z), \quad z \in \mathbb{C} \setminus (\Gamma \cup \Sigma_1 \cup [0, \infty)), \quad (3.13)$$

where $\log z$ is defined with a branch cut along $[0, \infty)$.

Proof. We note that

$$g'(z) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z-s} \frac{R_+(s)}{s} ds = \frac{1}{2} \frac{1}{2\pi i} \oint_{\gamma} \frac{1}{z-s} \frac{R(s)}{s} ds$$

where γ is a closed contour in $\mathbb{C} \setminus \Gamma$, that encircles Γ once in the clockwise direction, but does not encircle z and 0 . Then exactly as in the proof of Proposition 3.4, we deform the contour, and now pick up residues at z , 0 and ∞ . We find for $z \in \mathbb{C} \setminus \Gamma$,

$$\begin{aligned} g'(z) &= \frac{1}{2} \left[\operatorname{Res}_{s=z} \left(\frac{1}{z-s} \frac{R(s)}{s} \right) + \operatorname{Res}_{s=0} \left(\frac{1}{z-s} \frac{R(s)}{s} \right) - \operatorname{Res}_{s=\infty} \left(\frac{1}{z-s} \frac{R(s)}{s} \right) \right] \\ &= \frac{1}{2} \left[-\frac{R(z)}{z} + \frac{R(0)}{z} + 1 \right] \\ &= \frac{1}{2} \left[-\frac{R(z)}{z} + \frac{A}{z} + 1 \right]. \end{aligned} \quad (3.14)$$

Let $z \in \mathbb{C} \setminus (\Gamma \cup \Sigma_1 \cup [0, \infty))$. Integrating (3.14) from β to z along a curve in $\mathbb{C} \setminus (\Gamma \cup \Sigma_1 \cup [0, \infty))$, we find

$$g(z) = g(\beta) + \frac{1}{2}(A \log z + z) - \frac{1}{2}(A \log \beta + \beta) - \phi(z). \quad (3.15)$$

Thus (3.13) holds with $\ell = 2g(\beta) - (A \log \beta + \beta)$. $\square$

3.4 Jump properties of g

From Proposition 3.7 we obtain the following jump relations for g across the contour Σ . These jumps are crucial for the subsequent analysis.

Proposition 3.8 (a) *We have*

$$g_+(z) - g_-(z) = 2\pi i \quad \text{for } z \in \Sigma_1, \quad (3.16)$$

and

$$g_+(z) - g_-(z) = -\phi_+(z) + \phi_-(z) = -2\phi_+(z) = 2\phi_-(z) \quad \text{for } z \in \Gamma. \quad (3.17)$$

(b) *We have, with the same constant ℓ as in Proposition 3.7,*

$$g_+(z) + g_-(z) = A \log z + z + \ell \quad \text{for } z \in \Gamma, \quad (3.18)$$

$$g_+(z) + g_-(z) = A \log z + z + \ell - 2\tilde{\phi}(z) \quad \text{for } z \in \Sigma_1, \quad (3.19)$$

and

$$g_+(z) + g_-(z) = A \log z + z + \ell - 2\phi(z) \quad \text{for } z \in \Sigma_2. \quad (3.20)$$

Proof. In (3.13) we let z approach Σ , from the $+$ - and $-$ -sides, respectively, to obtain

$$g_{\pm}(z) = \frac{1}{2}(A \log z + z) + \frac{1}{2}\ell - \phi_{\pm}(z), \quad \text{for } z \in \Sigma. \quad (3.21)$$

Since ϕ changes sign across Γ , (3.17) and (3.18) immediately follow from (3.21). For $z \in \Sigma_1$, we have by (3.12) and (3.21)

$$\begin{aligned} g_+(z) - g_-(z) &= -\phi_+(z) + \phi_-(z) = -(\tilde{\phi}_+(z) - \pi i) + (\tilde{\phi}_-(z) + \pi i) \\ &= 2\pi i - \tilde{\phi}_+(z) + \tilde{\phi}_-(z), \end{aligned}$$

which is (3.16) as $\tilde{\phi}$ is analytic across Σ_1 . In addition, we have

$$\begin{aligned} g_+(z) + g_-(z) &= A \log z + z + \ell - \phi_+(z) - \phi_-(z) \\ &= A \log z + z + \ell - \tilde{\phi}_+(z) - \tilde{\phi}_-(z) \quad \text{for } z \in \Sigma_1. \end{aligned}$$

which yields (3.19). Similarly, (3.20) follows. $\square$

4 First two transformations: $Y \mapsto U \mapsto T$

4.1 First transformation $Y \mapsto U$

With the g -function and the constant ℓ from Proposition 3.7, we perform the first transformation of the Riemann–Hilbert problem.

Definition 4.1 We define for $z \in \mathbb{C} \setminus \Sigma$,

$$U(z) = e^{-n(\ell/2)\sigma_3} Y(z) e^{-ng(z)\sigma_3} e^{n(\ell/2)\sigma_3}. \quad (4.1)$$

Here, and in what follows, σ_3 denotes the Pauli matrix $\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, so that for example

$$e^{-ng(z)\sigma_3} = \begin{pmatrix} e^{-ng(z)} & 0 \\ 0 & e^{ng(z)} \end{pmatrix}.$$

From the Riemann–Hilbert problem for Y it follows by a straightforward calculation that U is the unique solution of the following Riemann–Hilbert problem.

Riemann–Hilbert problem for U :

The problem is to determine a 2×2 matrix valued function $U : \mathbb{C} \setminus \Sigma \rightarrow \mathbb{C}^{2 \times 2}$ such that

- (a) $U(z)$ is analytic for $z \in \mathbb{C} \setminus \Sigma$,
- (b) $U(z)$ possesses continuous boundary values for $z \in \Sigma$, denoted by $U_+(z)$ and $U_-(z)$, and

$$U_+(z) = U_-(z) \begin{pmatrix} e^{-n(g_+(z) - g_-(z))} & z^{-An} e^{-nz} e^{n(g_+(z) + g_-(z) - \ell)} \\ 0 & e^{n(g_+(z) - g_-(z))} \end{pmatrix} \quad (4.2)$$

for $z \in \Sigma$,

- (c) $U(z)$ behaves like the identity at infinity:

$$U(z) = I + O\left(\frac{1}{z}\right) \quad \text{as } z \rightarrow \infty, \quad z \in \mathbb{C} \setminus \Sigma. \quad (4.3)$$

The jump relation (4.2) for U has a different form on the three parts Γ , Σ_1 and Σ_2 . On Γ we have that $g_+ - g_- = -2\phi_+ = 2\phi_-$ by (3.17), and $g_+ + g_- = A \log z + z + \ell$ by (3.18) so that

$$U_+(z) = U_-(z) \begin{pmatrix} e^{2n\phi_+(z)} & 1 \\ 0 & e^{2n\phi_-(z)} \end{pmatrix} \quad \text{for } z \in \Gamma. \quad (4.4)$$

On Σ_1 we use (3.16) and (3.19) to obtain

$$U_+(z) = U_-(z) \begin{pmatrix} 1 & e^{-2n\tilde{\phi}(z)} \\ 0 & 1 \end{pmatrix} \quad \text{for } z \in \Sigma_1, \quad (4.5)$$

and similarly it follows that

$$U_+(z) = U_-(z) \begin{pmatrix} 1 & e^{-2n\phi(z)} \\ 0 & 1 \end{pmatrix} \quad \text{for } z \in \Sigma_2. \quad (4.6)$$

The transformation $Y \mapsto U$ has the effect of normalizing the Riemann–Hilbert problem at infinity. In addition, by the construction of Σ , we have that ϕ is real and positive on Σ_2 , and $\tilde{\phi}$ is real and positive on Σ_1 . So the jump matrices for U in (4.5) and (4.6) are close to the identity if n is large. Since ϕ has purely imaginary boundary values on both sides of Γ , the jump matrix for U on Γ in (4.4) has oscillatory diagonal entries.

4.2 Second transformation $U \mapsto T$

The jump matrix for U on Γ , see (4.4), factors as

$$\begin{pmatrix} e^{2n\phi_+(z)} & 1 \\ 0 & e^{2n\phi_-(z)} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ e^{2n\phi_-(z)} & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ e^{2n\phi_+(z)} & 1 \end{pmatrix}. \quad (4.7)$$

Observe that the first matrix in the right-hand side of (4.7) can be analytically continued to the $-$ -side of the contour Γ , and in doing so, the (1,2) entry becomes exponentially decaying in n . Similarly, the third matrix in the right-hand side of (4.7) can be analytically continued to the $+$ -side of the contour Γ , and in doing so, the (1,2) entry also becomes exponentially decaying in n . We are thus led to introduce the following “contour augmentation” step, as part of the steepest descent / stationary phase method for Riemann–Hilbert problems developed by Deift and Zhou. The oriented contour Σ^T consists of Σ plus two simple curves Σ_3 and Σ_4 from $\bar{\beta}$ to β , contained in Ω_+ and Ω_- , respectively, as shown in Figure 6. We choose Σ_3 and Σ_4 such that $\operatorname{Re} \phi(z) < 0$ on Σ_3 and Σ_4 .

It is possible to choose such curves. Indeed, ϕ is positive on Σ_2 , and its real part vanishes on Γ and on the other solid lines shown in Figure 4. So $\operatorname{Re} \phi > 0$ in the full region on the right. In the two other regions, bounded by the solid lines, we then have that $\operatorname{Re} \phi < 0$. Note that $\operatorname{Re} \phi$ does not change sign across Γ .

Note that by (3.12) we also have $\operatorname{Re} \tilde{\phi}(z) < 0$ on Σ_3 and Σ_4 .

Then $\mathbb{C} \setminus \Sigma^T$ has four connected components, denoted by Ω_1 , Ω_2 , Ω_3 , and Ω_4 as indicated in Figure 6.

Definition 4.2 We define $T : \mathbb{C} \setminus \Sigma^T \rightarrow \mathbb{C}^{2 \times 2}$ by

$$T(z) = U(z) \quad \text{for } z \in \Omega_1 \cup \Omega_4, \quad (4.8)$$




Figure 6: Contour $\Sigma^T = \Gamma \cup \bigcup_j \Sigma_j$ and the domains Ω_j , $j = 1, \dots, 4$, for the Riemann–Hilbert problem for T , for the value $A = 1.01$.

$$T(z) = U(z) \begin{pmatrix} 1 & 0 \\ -e^{2n\phi(z)} & 1 \end{pmatrix} \quad \text{for } z \in \Omega_2, \quad (4.9)$$

$$T(z) = U(z) \begin{pmatrix} 1 & 0 \\ e^{2n\phi(z)} & 1 \end{pmatrix} \quad \text{for } z \in \Omega_3, \quad (4.10)$$

Then from the Riemann–Hilbert problem for U and the factorization (4.7) we obtain that T is the unique solution of the following Riemann–Hilbert problem.

Riemann–Hilbert problem for T :

The problem is to determine a 2×2 matrix valued function $T : \mathbb{C} \setminus \Sigma^T \rightarrow \mathbb{C}^{2 \times 2}$ such that the following hold:

- (a) $T(z)$ is analytic for $z \in \mathbb{C} \setminus \Sigma^T$,
- (b) $T(z)$ possesses continuous boundary values for $z \in \Sigma^T$, denoted by $T_+(z)$ and $T_-(z)$,
and

$$T_+(z) = T_-(z) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad \text{for } z \in \Gamma, \quad (4.11)$$

$$T_+(z) = T_-(z) \begin{pmatrix} 1 & 0 \\ e^{2n\phi(z)} & 1 \end{pmatrix} \quad \text{for } z \in \Sigma_3 \cup \Sigma_4, \quad (4.12)$$

$$T_+(z) = T_-(z) \begin{pmatrix} 1 & e^{-2n\tilde{\phi}(z)} \\ 0 & 1 \end{pmatrix} \quad \text{for } z \in \Sigma_1, \quad (4.13)$$

and

$$T_+(z) = T_-(z) \begin{pmatrix} 1 & e^{-2n\phi(z)} \\ 0 & 1 \end{pmatrix} \quad \text{for } z \in \Sigma_2, \quad (4.14)$$

- (c) $T(z)$ behaves like the identity at infinity:

$$T(z) = I + O\left(\frac{1}{z}\right) \quad \text{as } z \rightarrow \infty, \quad z \in \mathbb{C} \setminus \Sigma^T. \quad (4.15)$$

5 Construction of the parametrix for T

5.1 Parametrix away from β and $\bar{\beta}$

As remarked following (4.7), the jump matrices appearing in (4.12) for $z \in \Sigma_3 \cup \Sigma_4$ are exponentially close to the identity matrix away from β and $\bar{\beta}$. Similarly, the jump matrices appearing in (4.13) and (4.14) are also exponentially close to the identity matrix away from β and $\bar{\beta}$. This hints that these portions of the contour on which the Riemann–Hilbert problem for T is posed should be somehow negligible. Thus we expect that the leading order asymptotics is determined by the solution N of the following model Riemann–Hilbert problem:

Riemann–Hilbert problem for N :

The problem is to determine $N : \mathbb{C} \setminus \Gamma \rightarrow \mathbb{C}^{2 \times 2}$ such that the following hold.

- (a) $N(z)$ is analytic for $z \in \mathbb{C} \setminus \Gamma$,
- (b) $N(z)$ possesses continuous boundary values for $z \in \Gamma \setminus \{\beta, \bar{\beta}\}$, denoted by $N_+(z)$ and $N_-(z)$, and

$$N_+(z) = N_-(z) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad \text{for } z \in \Gamma \setminus \{\beta, \bar{\beta}\}, \quad (5.1)$$

- (c) $N(z) = I + O\left(\frac{1}{z}\right)$ for $z \rightarrow \infty$.

This Riemann–Hilbert problem for N is solved explicitly by, see [10, p.1520] or [6, p.200],

$$N(z) = \begin{pmatrix} \frac{a(z)+a(z)^{-1}}{2} & \frac{a(z)-a(z)^{-1}}{2i} \\ \frac{a(z)-a(z)^{-1}}{-2i} & \frac{a(z)+a(z)^{-1}}{2} \end{pmatrix}, \quad (5.2)$$

where

$$a(z) = \frac{(z - \beta)^{1/4}}{(z - \bar{\beta})^{1/4}}. \quad (5.3)$$

The branches of the roots in (5.3) are chosen such that $a(z)$ is analytic on $\mathbb{C} \setminus \Gamma$ and $\lim_{z \rightarrow \infty} a(z) = 1$.

Remark 5.1 The solution (5.2) is not the only solution to the Riemann–Hilbert problem for N . It is the unique solution that satisfies, in addition to (a), (b), and (c), the condition

- (d) Near the endpoints β and $\bar{\beta}$, we have

$$N(z) = O(|z - \beta|^{-1/4}) \quad \text{as } z \rightarrow \beta,$$

$$N(z) = O(|z - \bar{\beta}|^{-1/4}) \quad \text{as } z \rightarrow \bar{\beta},$$

with the O -term being taken entry-wise.

Remark 5.2 For explicit calculations, it is useful to have an alternative expression for N . For $z = x$ real, it is clear from (5.3) that $a(x)$ has modulus one. If $\arg(a(x)) = \theta(x)$, then (5.2) shows

$$N(x) = \begin{pmatrix} \cos \theta(x) & \sin \theta(x) \\ -\sin \theta(x) & \cos \theta(x) \end{pmatrix}.$$

Since $\tan(2\theta(x)) = -\frac{2\sqrt{A-1}}{x-2+A}$, we then find

$$N(z) = \begin{pmatrix} \cos(\frac{1}{2} \arctan(\frac{2\sqrt{A-1}}{z-2+A})) & -\sin(\frac{1}{2} \arctan(\frac{2\sqrt{A-1}}{z-2+A})) \\ \sin(\frac{1}{2} \arctan(\frac{2\sqrt{A-1}}{z-2+A})) & \cos(\frac{1}{2} \arctan(\frac{2\sqrt{A-1}}{z-2+A})) \end{pmatrix} \quad (5.4)$$

first for $z = x$ real, but then also for arbitrary $z \in \mathbb{C} \setminus \Gamma$ by analytic continuation. We have to take the appropriate branch of the multivalued arctan function in (5.4). Using trigonometric identities, one may then check that (5.4) reduces to

$$N(z) = \begin{pmatrix} \left(\frac{1+R'(z)}{2}\right)^{1/2} & -\left(\frac{1-R'(z)}{2}\right)^{1/2} \\ \left(\frac{1-R'(z)}{2}\right)^{1/2} & \left(\frac{1+R'(z)}{2}\right)^{1/2} \end{pmatrix} \quad \text{for } z \in \mathbb{C} \setminus \Gamma, \quad (5.5)$$

where as usual we have $R(z) = (z - \beta)^{1/2}(z - \bar{\beta})^{1/2}$. We can also verify directly that (5.5) solves the Riemann–Hilbert problem for N .

5.2 Parametrix near β

The next step is a local analysis around the points β and $\bar{\beta}$. We need to construct a local parametrix P in a neighborhood $U_\delta = \{z \in \mathbb{C} \mid |z - \beta| < \delta\}$ of β such that

- P satisfies the jumps for T exactly in U_δ ,
- P matches N on the boundary of U_δ up to order $1/n$.

See Figure 7 for the contours $\Sigma^T \cap U_\delta$.

More precisely, we have

Riemann–Hilbert problem for P :

The problem is to determine, for a given $\delta > 0$ sufficiently small, a 2×2 matrix valued function $P : \overline{U}_\delta \setminus \Sigma^T \rightarrow \mathbb{C}^{2 \times 2}$ such that

- $P(z)$ is analytic for $z \in U_\delta \setminus \Sigma^T$, and continuous on $\overline{U}_\delta \setminus \Sigma^T$,
- $P(z)$ possesses continuous boundary values for $z \in \Sigma^T \cap U_\delta$, denoted by $P_+(z)$ and $P_-(z)$, and

$$P_+(z) = P_-(z) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad \text{for } z \in \Gamma \cap U_\delta, \quad (5.6)$$

$$P_+(z) = P_-(z) \begin{pmatrix} 1 & 0 \\ e^{2n\phi(z)} & 1 \end{pmatrix} \quad \text{for } z \in (\Sigma_3 \cup \Sigma_4) \cap U_\delta, \quad (5.7)$$

and

$$P_+(z) = P_-(z) \begin{pmatrix} 1 & e^{-2n\phi(z)} \\ 0 & 1 \end{pmatrix} \quad \text{for } z \in \Sigma_2 \cap U_\delta, \quad (5.8)$$




Figure 7: Neighborhood U_δ of β and the parts of the contours Γ , Σ_2 , Σ_3 , and Σ_4 that are within U_δ . The value of A is 1.01.

(c) There exists a constant $C > 0$ such that for every $z \in \partial U_\delta \setminus \Sigma^T$,

$$\|P(z)N^{-1}(z) - I\| \leq \frac{C}{n}, \quad (5.9)$$

where $\|\cdot\|$ is any matrix norm.

The construction of P follows along the same lines as given by Deift et al. [10]. From its definition (3.10) it is easy to see that the ϕ -function has a convergent expansion

$$\phi(z) = (z - \beta)^{3/2} \sum_{k=0}^{\infty} c_k (z - \beta)^k, \quad c_0 \neq 0, \quad (5.10)$$

in a neighborhood of β . The factor $(z - \beta)^{3/2}$ is defined with a cut along $\Gamma \cup \Sigma_1$. Then

$$f(z) = \left[\frac{3}{2} \phi(z) \right]^{2/3} \quad (5.11)$$

is defined and analytic in a neighborhood of β . We choose the $2/3$ -root with a cut along Γ and such that $f(z) > 0$ for $z \in \Sigma_2$. Recall $\phi > 0$ on Σ_2 .

Then $f(\beta) = 0$ and $f'(\beta) \neq 0$. Therefore we can and do choose δ so small that $\zeta = f(z)$ is a one-to-one mapping from U_δ onto a convex neighborhood $f(U_\delta)$ of $\zeta = 0$. Under the mapping $\zeta = f(z)$, we then have that $\Sigma_2 \cap U_\delta$ corresponds to $(0, \infty) \cap f(U_\delta)$ and that $\Gamma \cap U_\delta$ corresponds to $(-\infty, 0] \cap f(U_\delta)$.

Now we specify how to choose Σ^T near β . For an arbitrary, but fixed $\sigma \in (\pi/3, \pi)$, we choose Σ_3 and Σ_4 such that f maps $\Sigma_3 \cap U_\delta$ and $\Sigma_4 \cap U_\delta$ onto $\{\zeta \in f(U_\delta) \mid \arg \zeta = \sigma\}$ and $\{\zeta \in f(U_\delta) \mid \arg \zeta = -\sigma\}$, respectively.

Proposition 5.3 *The Riemann–Hilbert problem for P is solved by*

$$P(z) = E(z) \Psi^\sigma (n^{2/3} f(z)) e^{n\phi(z)\sigma_3} \quad (5.12)$$

where

$$E(z) = \sqrt{\pi} e^{\frac{\pi i}{6}} \begin{pmatrix} 1 & -1 \\ -i & -i \end{pmatrix} \left(\frac{n^{1/6} f(z)^{1/4}}{a(z)} \right)^{\sigma_3} \quad (5.13)$$

and Ψ^σ is an explicit matrix valued function built out of the Airy function Ai and its derivative Ai' as follows

$$\Psi^\sigma(\zeta) = \begin{cases} \begin{pmatrix} \text{Ai}(\zeta) & \text{Ai}(\omega^2 \zeta) \\ \text{Ai}'(\zeta) & \omega^2 \text{Ai}'(\omega^2 \zeta) \end{pmatrix} e^{-\frac{\pi i}{6} \sigma_3} & \text{for } 0 < \arg \zeta < \sigma, \\ \begin{pmatrix} \text{Ai}(\zeta) & \text{Ai}(\omega^2 \zeta) \\ \text{Ai}'(\zeta) & \omega^2 \text{Ai}'(\omega^2 \zeta) \end{pmatrix} e^{-\frac{\pi i}{6} \sigma_3} \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix} & \text{for } \sigma < \arg \zeta < \pi, \\ \begin{pmatrix} \text{Ai}(\zeta) & -\omega^2 \text{Ai}(\omega \zeta) \\ \text{Ai}'(\zeta) & -\text{Ai}'(\omega \zeta) \end{pmatrix} e^{-\frac{\pi i}{6} \sigma_3} \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} & \text{for } -\pi < \arg \zeta < -\sigma, \\ \begin{pmatrix} \text{Ai}(\zeta) & -\omega^2 \text{Ai}(\omega \zeta) \\ \text{Ai}'(\zeta) & -\text{Ai}'(\omega \zeta) \end{pmatrix} e^{-\frac{\pi i}{6} \sigma_3} & \text{for } -\sigma < \arg \zeta < 0, \end{cases} \quad (5.14)$$

with $\omega = e^{2\pi i/3}$.

Proof. The proof is similar to [10, p.1523–1525]. $\square$

Remark 5.4 An analysis of the proof in [10] shows that the constant C in (5.9) can be taken uniformly for σ in a compact subset K_σ of $(\pi/3, \pi)$, for A in a compact subset K_A of $(1, \infty)$, and for δ uniformly in some interval (δ_0, δ_1) depending on K_σ and K_A .

A similar remark applies to the parametrix $\tilde{P}$ to be constructed below.

5.3 Parametrix near $\bar{\beta}$

A similar construction yields a parametrix $\tilde{P}$ in a neighborhood $\tilde{U}_\delta = \{z \mid |z - \bar{\beta}| < \delta\}$ that satisfies the following Riemann–Hilbert problem.

Riemann–Hilbert problem for $\tilde{P}$:

The problem is to determine, for a given $\delta > 0$ sufficiently small, a 2×2 matrix valued function $\tilde{P} : \tilde{U}_\delta \setminus \Sigma^T \rightarrow \mathbb{C}^{2 \times 2}$ such that

- (a) $\tilde{P}(z)$ is analytic for $z \in U_\delta \setminus \Sigma^T$, and continuous on $\tilde{U}_\delta \setminus \Sigma^T$,
- (b) $\tilde{P}(z)$ possesses continuous boundary values for $z \in \Sigma^T \cap \tilde{U}_\delta$, denoted by $\tilde{P}_+(z)$ and $\tilde{P}_-(z)$, and

$$\tilde{P}_+(z) = \tilde{P}_-(z) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad \text{for } z \in \Gamma \cap \tilde{U}_\delta, \quad (5.15)$$

$$\tilde{P}_+(z) = \tilde{P}_-(z) \begin{pmatrix} 1 & 0 \\ e^{2n\tilde{\phi}(z)} & 1 \end{pmatrix} \quad \text{for } z \in (\Sigma_3 \cup \Sigma_4) \cap \tilde{U}_\delta, \quad (5.16)$$

and

$$\tilde{P}_+(z) = \tilde{P}_-(z) \begin{pmatrix} 1 & e^{-2n\tilde{\phi}(z)} \\ 0 & 1 \end{pmatrix} \quad \text{for } z \in \Sigma_1 \cap \tilde{U}_\delta. \quad (5.17)$$

Recall that the $\tilde{\phi}$ -function is defined in (3.11).

(c) There exists a constant $C > 0$ such that for every $z \in \partial\tilde{U}_\delta \setminus \Sigma^T$,

$$\|\tilde{P}(z)N^{-1}(z) - I\| \leq \frac{C}{n}. \quad (5.18)$$

There is a one-to-one analytic mapping $\zeta = \tilde{f}(z)$ such that

$$\frac{2}{3}(-\tilde{f}(z))^{3/2} = \tilde{\phi}(z). \quad (5.19)$$

Then $\tilde{f}$ maps $\tilde{U}_\delta$ onto a convex neighborhood of $\zeta = 0$.

Proposition 5.5 *The Riemann–Hilbert problem for $\tilde{P}$ is solved by*

$$\tilde{P}(z) = \tilde{E}(z)\tilde{\Psi}^\sigma(n^{2/3}\tilde{f}(z))e^{n\tilde{\phi}(z)\sigma_3} \quad (5.20)$$

with

$$\tilde{E}(z) = \sqrt{\pi}e^{\pi i/6} \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix} \left(n^{1/6}(-\tilde{f}(z))^{1/4}a(z) \right)^{\sigma_3}, \quad (5.21)$$

and

$$\tilde{\Psi}^\sigma(\zeta) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \Psi^\sigma(-\zeta) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (5.22)$$

Proof. This follows as in [10, p.1527]. $\square$

Remark 5.6 As in [10] there is a full asymptotic expansion of PN^{-1} and $\tilde{P}N^{-1}$ in inverse powers of n . This expansion leads to uniform asymptotic expansions for the generalized Laguerre polynomials. In the present paper we will compute asymptotics for the polynomials to first order, but we will not carry out the further computations to determine the coefficients of a complete asymptotic expansion.

6 Final transformation $T \mapsto S$

Using N , P , and $\tilde{P}$, we define for every $n \in \mathbb{N}$,

$$S(z) = T(z)N(z)^{-1} \quad \text{for } z \in \mathbb{C} \setminus (\Sigma^T \cup \overline{U}_\delta \cup \overline{\tilde{U}}_\delta), \quad (6.1)$$

$$S(z) = T(z)P(z)^{-1} \quad \text{for } z \in U_\delta \setminus \Sigma^T, \quad (6.2)$$

$$S(z) = T(z)\tilde{P}(z)^{-1} \quad \text{for } z \in \tilde{U}_\delta \setminus \Sigma^T. \quad (6.3)$$

Then S is defined and analytic on $\mathbb{C} \setminus (\Sigma^T \cup \partial U_\delta \cup \partial \tilde{U}_\delta)$. However it follows from the construction that S has no jumps on Γ and on $\Sigma^T \cap (U_\delta \cup \tilde{U}_\delta)$. Therefore S has an analytic continuation to $\mathbb{C} \setminus \Sigma^S$ (also denoted by S), where Σ^S is the contour indicated in Figure 8 for $A = 1.01$.

Formally, the contours Σ_j^S are given by $\Sigma_j^S = \Sigma_j \setminus (\overline{U}_\delta \cup \overline{\tilde{U}}_\delta)$ for $j = 1, 2, 3, 4$, and the domains Ω_j^S are given by $\Omega_1^S = \Omega_1 \setminus (\overline{U}_\delta \cup \overline{\tilde{U}}_\delta)$, $\Omega_2^S = (\Omega_2 \cup \Gamma \cup \Omega_3) \setminus (\overline{U}_\delta \cup \overline{\tilde{U}}_\delta)$, and $\Omega_3^S = \Omega_4 \setminus (\overline{U}_\delta \cup \overline{\tilde{U}}_\delta)$.

Then S satisfies the following Riemann–Hilbert problem.




Figure 8: Contour $\Sigma^S = \partial U_\delta \cup \partial \tilde{U}_\delta \cup \bigcup_j \Sigma_j^S$ and the domains U_δ , $\tilde{U}_\delta$, and Ω_j^S , $j = 1, 2, 3$, for the Riemann–Hilbert problem for S , for the value $A = 1.01$.

Riemann–Hilbert problem for S :

The problem is to determine $S : \mathbb{C} \setminus \Sigma^S \rightarrow \mathbb{C}^{2 \times 2}$ such that the following hold.

- (a) $S(z)$ is analytic for $z \in \mathbb{C} \setminus \Sigma^S$,
- (b) $S(z)$ possesses continuous boundary values for $z \in \Sigma^S$, denoted by $S_+(z)$ and $S_-(z)$, and

$$S_+(z) = S_-(z)P(z)N(z)^{-1} \quad \text{for } z \in \partial U_\delta, \quad (6.4)$$

$$S_+(z) = S_-(z)\tilde{P}(z)N(z)^{-1} \quad \text{for } z \in \partial \tilde{U}_\delta, \quad (6.5)$$

$$S_+(z) = S_-(z)N(z) \begin{pmatrix} 1 & 0 \\ e^{2n\phi(z)} & 1 \end{pmatrix} N(z)^{-1} \quad \text{for } z \in \Sigma_3^S \cup \Sigma_4^S, \quad (6.6)$$

$$S_+(z) = S_-(z)N(z) \begin{pmatrix} 1 & e^{-2n\tilde{\phi}(z)} \\ 0 & 1 \end{pmatrix} N(z)^{-1} \quad \text{for } z \in \Sigma_1^S, \quad (6.7)$$

$$S_+(z) = S_-(z)N(z) \begin{pmatrix} 1 & e^{-2n\phi(z)} \\ 0 & 1 \end{pmatrix} N(z)^{-1} \quad \text{for } z \in \Sigma_2^S, \quad (6.8)$$

- (c) $S(z) = I + O\left(\frac{1}{z}\right)$ for $z \rightarrow \infty$.

The jump matrices for S are close to the identity matrix if n is large. Indeed, by (5.9) and (5.18) we have

$$\|P(z)N^{-1}(z) - I\| \leq \frac{C}{n} \quad \text{for } z \in \partial U_\delta,$$

and

$$\|\tilde{P}(z)N^{-1}(z) - I\| \leq \frac{C}{n} \quad \text{for } z \in \partial\tilde{U}_\delta,$$

with a constant C that is independent of z . The constant can also be chosen independently of the value of A for A in a compact subset of $(1, \infty)$, see Remark 5.4.

The jump matrices in (6.6)–(6.8) are exponentially close to the identity matrix.

7 Asymptotic for polynomials $L_n^{(\alpha_n)}(nz)$

7.1 Asymptotics for S

In Sections 3–6 the values of n and α were assumed to be fixed. In order to study the asymptotics, we now let $\alpha = \alpha_n$ depend on n and we write $A_n = -\frac{\alpha_n}{n}$. We assume that $A_n > 1$ for every n , and

$$\lim_{n \rightarrow \infty} -\frac{\alpha_n}{n} = \lim_{n \rightarrow \infty} A_n = A > 1. \quad (7.1)$$

Because of the n -dependence of A_n , all of the notions and results introduced in Sections 3–6 are n -dependent. For example, we have that the curves Γ , Σ , Σ^T , and Σ^S are all varying with n , and so we denote them by Γ_n , Σ_n , Σ_n^T , and Σ_n^S . Likewise, we have that the functions R , g , ϕ , and $\tilde{\phi}$, as well as all matrix-valued functions are n -dependent, and we also use a subscript n to denote their dependence on n . If now we use Γ , Σ , R , g , etc., without subscript n , then this refers to the limiting case. Due to (7.1) we have that the curves Γ_n tend to the limiting curve Γ , etc.

At the end of the previous section, we observed that the jump matrix for S_n is $I + O(1/n)$ uniformly on Σ_n^S as $n \rightarrow \infty$. In addition, the jump matrix converges to the identity matrix as $z \rightarrow \infty$ along the unbounded components of Σ_n^S sufficiently fast, so that the jump matrix is also close to I in the L^2 -sense. Since the contours Σ_n^S are only slightly varying with n , we may follow arguments as those given in [2] and [6], to conclude that

$$S_n(z) = I + O\left(\frac{1}{n}\right) \quad \text{as } n \rightarrow \infty \quad (7.2)$$

uniformly for $z \in \mathbb{C} \setminus \Sigma_n^S$. Briefly, the Riemann–Hilbert problem for S_n is equivalent to a system of singular integral equations. This system is of the form $\gamma(\zeta) - F(\gamma)(\zeta) = g(\zeta)$, where γ is a matrix valued function defined on the contour Σ_n^S , and where the singular integral operator $F(\gamma)(\zeta)$ has operator norm $O(\frac{1}{n})$. The system of singular integral equations can therefore be solved by Neumann series in powers of $\frac{1}{n}$, and this leads to a series expansion for the solution of the Riemann–Hilbert problem.

Using more precise information on the jump matrix for S_n as indicated in Remark 5.6, and with additional assumptions on the limit (7.1), one is able to obtain a full asymptotic expansion for $S_n(z)$ in powers of $1/n$. This in return would give a full asymptotic expansion for the generalized Laguerre polynomials.

7.2 Strong asymptotics for generalized Laguerre polynomials

Unraveling the steps $Y_n \mapsto U_n \mapsto T_n \mapsto S_n$ and using (7.2), we obtain strong asymptotics for Y_n in all regions of the complex plane. In particular we are interested in the $(1,1)$ entry of Y_n , since this is the monic generalized Laguerre polynomial.

Theorem 7.1 Suppose for each $n \in \mathbb{N}$, we have a parameter $\alpha_n < -n$ such that (7.1) holds where $A_n = -\frac{\alpha_n}{n}$. Then we have the following asymptotic results for the generalized Laguerre polynomials $L_n^{(\alpha_n)}(nz)$ as $n \rightarrow \infty$.

(a) **(asymptotics away from Γ)**

Uniformly for z in compact subsets of $\mathbb{C} \setminus \Gamma$, we have as $n \rightarrow \infty$,

$$L_n^{(\alpha_n)}(nz) = \frac{(-n)^n}{n!} e^{ng_n(z)} \left(\frac{1 + R'_n(z)}{2} \right)^{1/2} \left(1 + O\left(\frac{1}{n}\right) \right). \quad (7.3)$$

(b) **(asymptotics on +side of Γ_n , away from endpoints)**

Uniformly for z on the +side of Γ_n away from β and $\bar{\beta}$, we have as $n \rightarrow \infty$,

$$\begin{aligned} L_n^{(\alpha_n)}(nz) &= \frac{(-n)^n}{n!} e^{ng_n(z)} \left(\frac{1 + R'_n(z)}{2} \right)^{1/2} \\ &\times \left[1 - \left(\frac{1 - R'_n(z)}{1 + R'_n(z)} \right)^{1/2} e^{2n\phi_n(z)} + O\left(\frac{1}{n}\right) \right]. \end{aligned} \quad (7.4)$$

(c) **(asymptotics on --side of Γ_n , away from endpoints)**

Uniformly for z on the --side of Γ_n away from β and $\bar{\beta}$, we have as $n \rightarrow \infty$,

$$\begin{aligned} L_n^{(\alpha_n)}(nz) &= \frac{(-n)^n}{n!} e^{ng_n(z)} \left(\frac{1 + R'_n(z)}{2} \right)^{1/2} \\ &\times \left[1 + \left(\frac{1 - R'_n(z)}{1 + R'_n(z)} \right)^{1/2} e^{2n\phi_n(z)} + O\left(\frac{1}{n}\right) \right]. \end{aligned} \quad (7.5)$$

(d) **(asymptotics near β)**

Uniformly for z in a (small) neighborhood of β , we have as $n \rightarrow \infty$,

$$\begin{aligned} L_n^{(\alpha_n)}(nz) &= \frac{(-n)^n}{n!} \exp\left(\frac{n}{2}(A_n \log z + z + \ell)\right) \sqrt{\pi} \\ &\times \left[\left(\frac{z - \bar{\beta}_n}{z - \beta_n} \right)^{1/4} (n^{2/3} f_n(z))^{1/4} \text{Ai}(n^{2/3} f_n(z)) \left(1 + O\left(\frac{1}{n}\right) \right) \right. \\ &\quad \left. - \left(\frac{z - \beta_n}{z - \bar{\beta}_n} \right)^{1/4} (n^{2/3} f_n(z))^{-1/4} \text{Ai}'(n^{2/3} f_n(z)) \left(1 + O\left(\frac{1}{n}\right) \right) \right]. \end{aligned} \quad (7.6)$$

Proof. (a) Let K be a compact subset of $\mathbb{C} \setminus \Gamma$. Then, for n large enough, say $n \geq n_0$, we have $K \subset \mathbb{C} \setminus \Gamma_n$. Let $n \geq n_0$. While choosing the contours $(\Sigma_3)_n$ and $(\Sigma_4)_n$ in Section 4, we then take care that $(\Omega_2)_n$ and $(\Omega_3)_n$ do not meet the compact K , see Figure 6. We also choose δ so that the disks of radius δ around β_n and $\bar{\beta}_n$ are disjoint from K . Then $K \subset (\Omega_1^S)_n \cup (\Omega_3^S)_n$, see Figure 8. For $z \in K$, we then have by (4.1), (4.8), and (6.1)

$$\begin{aligned} Y_n(z) &= e^{n(\ell_n/2)\sigma_3} U_n(z) e^{ng_n(z)\sigma_3} e^{-n(\ell_n/2)\sigma_3} \\ &= e^{n(\ell_n/2)\sigma_3} T_n(z) e^{ng_n(z)\sigma_3} e^{-n(\ell_n/2)\sigma_3} \\ &= e^{n(\ell_n/2)\sigma_3} S_n(z) N_n(z) e^{ng_n(z)\sigma_3} e^{-n(\ell_n/2)\sigma_3}. \end{aligned} \quad (7.7)$$

Using (2.8), (5.5), and (7.2), and the fact that $|(N_n)_{11}(z)|$ is bounded away from zero on K , we obtain (7.3) from (7.7).

(b) For $z \in (\Omega_2)_n \setminus ((U_\delta)_n \cup (\tilde{U}_\delta)_n)$, we have by (4.1) and (4.9)

$$(Y_n)_{11}(z) = (U_n)_{11}(z)e^{ng_n(z)} = [(T_n)_{11}(z) + (T_n)_{12}(z)e^{2n\phi_n(z)}]e^{ng_n(z)} \quad (7.8)$$

Since $T_n = S_n N_n$ by (6.1), $S_n = I + O(1/n)$, and the entries of N_n are uniformly bounded away from zero in the region under consideration, we then get

$$(Y_n)_{11}(z) = [(N_n)_{11}(z) + (N_n)_{12}(z)e^{2n\phi_n(z)} + O(1/n)]e^{ng_n(z)}, \quad (7.9)$$

uniformly for $z \in (\Omega_2)_n \setminus ((U_\delta)_n \cup (\tilde{U}_\delta)_n)$. This proves (7.4).

(c) This is proved similarly as part (b). The only difference is that for $z \in (\Omega_3)_n \setminus ((U_\delta)_n \cup (\tilde{U}_\delta)_n)$ we use (4.10) instead of (4.9). This leads to

$$(Y_n)_{11}(z) = [(T_n)_{11}(z) - (T_n)_{12}(z)e^{2n\phi_n(z)}]e^{ng_n(z)}$$

instead of (7.8). The rest of the proof is the same.

(d) The proof is as in [10, p.1539]. Let U_ϵ be an ϵ -neighborhood of β with $\epsilon < \delta$. Then $U_\epsilon \subset (U_\delta)_n$ for all large enough n , say $n \geq n_1$. Let $z \in U_\epsilon$ and $n \geq n_1$. For $z \in (\Omega_1)_n$, we find by (4.1), (4.8), and (6.2) that

$$\begin{aligned} (Y_n)_{11}(z) &= (U_n)_{11}(z)e^{ng_n(z)} = (T_n)_{11}(z)e^{ng_n(z)} \\ &= [(S_n)_{11}(z)(P_n)_{11}(z) + (S_n)_{12}(z)(P_n)_{21}(z)]e^{ng_n(z)}. \end{aligned} \quad (7.10)$$

By Proposition 5.3 we have

$$P_n(z) = E_n(z)\Psi^\sigma(\zeta)e^{n\phi_n(z)\sigma_3}$$

where $\zeta = n^{2/3}f_n(z)$. For $z \in (\Omega_1)_n$, we have that ζ belongs to the sector $0 < \arg \zeta < \sigma$, so that the first formula for $\Psi^\sigma(\zeta)$ in (5.14) applies. It follows that

$$\begin{pmatrix} (P_n)_{11}(z) \\ (P_n)_{12}(z) \end{pmatrix} = E_n(z) \begin{pmatrix} \text{Ai}(\zeta) \\ \text{Ai}'(\zeta) \end{pmatrix} e^{-\frac{\pi i}{6}} e^{n\phi_n(z)}. \quad (7.11)$$

Using the definition (5.13) of $E_n(z)$, we obtain from (7.11)

$$\begin{pmatrix} (P_n)_{11}(z) \\ (P_n)_{12}(z) \end{pmatrix} = \sqrt{\pi} \begin{pmatrix} 1 & -1 \\ -i & -i \end{pmatrix} \begin{pmatrix} \zeta^{1/4} \\ \frac{1}{a_n(z)} \end{pmatrix} \begin{pmatrix} \text{Ai}(\zeta) \\ \text{Ai}'(\zeta) \end{pmatrix} e^{n\phi_n(z)}.$$

Then

$$\begin{aligned} &(S_n)_{11}(z)(P_n)_{11}(z) + (S_n)_{12}(z)(P_n)_{21}(z) \\ &= \sqrt{\pi} \begin{pmatrix} 1 + O(\frac{1}{n}) & -1 + O(\frac{1}{n}) \end{pmatrix} \begin{pmatrix} \frac{\zeta^{1/4}}{a_n(z)} \text{Ai}(\zeta) \\ \frac{a_n(z)}{\zeta^{1/4}} \text{Ai}'(\zeta) \end{pmatrix} e^{n\phi_n(z)}. \end{aligned} \quad (7.12)$$

Combining (7.10) and (7.12) with $\zeta = n^{2/3}f_n(z)$, we obtain an expression for $(Y_n)_{11}(z)$, which leads to (7.6) in view of (2.4) and (3.13).

Next, for z in the other regions $(\Omega_j)_n$, $j = 2, 3, 4$, similar calculations lead to the same expression for $(Y_{11})_n(z)$. Hence (7.6) holds uniformly for $z \in U_\epsilon$.

This completes the proof of Theorem 7.1. $\square$

Remark 7.2 Since α_n is real, we have $L_n^{(\alpha_n)}(n\bar{z}) = \overline{L_n^{(\alpha_n)}(nz)}$, and so (7.6) also describes the asymptotic behavior near $\bar{\beta}$.

7.3 Asymptotics for zeros

From Theorem 7.1 we deduce the following results concerning the zeros of the generalized Laguerre polynomials $L_n^{(\alpha_n)}(nz)$.

Corollary 7.3 *Suppose we are in the same situation as in Theorem 7.1.*

- (a) **(All zeros tend to Γ)** *For every neighborhood Ω of Γ , there is n_0 such that for every $n \geq n_0$, all zeros of $L_n^{(\alpha_n)}(nz)$ are in Ω .*
- (b) **(Zeros are on the --side)** *For every $\delta > 0$, there is n_0 such that for every $n \geq n_0$, there are no zeros of $L_n^{(\alpha_n)}(nz)$ in the region $(\Omega_2)_n \setminus ((U_\delta) \cup (\tilde{U}_\delta))$.*

Proof. (a) This is immediate from the asymptotic formula (7.3) since $1 + R'_n(z) \neq 0$.

(b) Suppose z is a zero of $L_n^{(\alpha_n)}(nz)$ lying in $(\Omega_2)_n \setminus ((U_\delta) \cup (\tilde{U}_\delta))$, that is, on the +side of Γ_n . Then we have by (7.4)

$$\left(\frac{1 - R'_n(z)}{1 + R'_n(z)} \right)^{1/2} e^{2n\phi_n(z)} = 1 + O\left(\frac{1}{n}\right). \quad (7.13)$$

We know that $\operatorname{Re} \phi_n(z) \leq 0$. In order to obtain a contradiction it is thus enough to enough to prove that

$$|1 - R'_n(z)| < |1 + R'_n(z)|. \quad (7.14)$$

Equality holds in (7.14) if and only if $R'_n(z)$ is purely imaginary, so that $(R'_n)^2(z)$ is real and negative. From the definition (3.2) of R_n , we get

$$(R'_n)^2(z) = \frac{(z - \frac{\beta_n + \bar{\beta}_n}{2})^2}{(z - \beta_n)(z - \bar{\beta}_n)} = \frac{(z - \operatorname{Re} \beta_n)^2}{(z - \operatorname{Re} \beta_n)^2 + (\operatorname{Im} \beta_n)^2}.$$

This is real and negative if and only if z belongs to the vertical segment connecting $\bar{\beta}_n$ and β_n . Consequently, this is the set where $|1 - R'_n(z)| = |1 + R'_n(z)|$. The vertical segment and Γ form a closed contour, and it may be checked that (7.14) holds for z outside of this contour, which includes the region $(\Omega_2)_n \setminus ((U_\delta) \cup (\tilde{U}_\delta))$.

So we have a contradiction, and it follows that there are no zeros in $(\Omega_2)_n \setminus ((U_\delta) \cup (\tilde{U}_\delta))$ for n large enough. $\square$

Remark 7.4 From the uniform asymptotics (7.6) in a neighborhood of β it is possible to derive asymptotics for the extreme zeros of $L_n^{(\alpha_n)}(nz)$. For example, it follows that for fixed $k \in \mathbb{N}$ and $n \geq k$, there is a zero $z_{k,n}$ of $L_n^{(\alpha_n)}(nz)$ such that

$$z_{k,n} = \beta_n - \frac{\iota_k}{f'_n(\beta_n)n^{2/3}} + O\left(\frac{1}{n}\right), \quad (7.15)$$

where $-\iota_k$ is the k th largest (negative) zero of the Airy function $\operatorname{Ai}(x)$.

References

- [1] M. Abramowitz and I.A. Stegun, Handbook of mathematical functions, Dover Publications, New York, 1966.
- [2] J. Baik, P. Deift, and K. Johansson, On the distribution of the length of the longest increasing subsequence of random permutations, *J. Amer. Math. Soc.* 12 (1999), 1119–1178.
- [3] P. Bleher and A. Its, Semiclassical asymptotics of orthogonal polynomials, Riemann–Hilbert problem, and universality in the matrix model, *Ann. Math.* 150 (1999), 185–266.
- [4] M. de Bruin, E.B. Saff, and R.S. Varga, On the zeros of generalized Bessel polynomials, I and II, *Indag. Math.* 43 (1981), 1–25.
- [5] A.J. Carpenter, Asymptotics for the zeros of the generalized Bessel polynomials, *Numer. Math.* 62 (1992), 465–482.
- [6] P. Deift, Orthogonal polynomials and random matrices: a Riemann–Hilbert approach, Courant Lecture Notes 3, Courant Institute 1999.
- [7] P. Deift, Integrable systems and combinatorial theory, *Notices Amer. Math. Soc.* 47 (2000), 631–640.
- [8] P. Deift, A. Its, and X. Zhou, A Riemann–Hilbert approach to asymptotic problems arising in the theory of random matrix models, and also in the theory of integrable statistical mechanics, *Ann. Math.* 146 (1997), 149–235.
- [9] P. Deift, T. Kriecherbauer, K.T-R McLaughlin, S. Venakides, and X. Zhou, Uniform asymptotics for polynomials orthogonal with respect to varying exponential weights and applications to universality questions in random matrix theory, *Comm. Pure Appl. Math.* 52 (1999), 1335–1425.
- [10] P. Deift, T. Kriecherbauer, K.T-R McLaughlin, S. Venakides, and X. Zhou, Strong asymptotics of orthogonal polynomials with respect to exponential weights, *Comm. Pure Appl. Math.* 52 (1999), 1491–1552.
- [11] P. Deift, S. Venakides, and X. Zhou, New results in small dispersion KdV by an extension of the steepest descent method for Riemann–Hilbert problems. *Internat. Math. Res. Notices* 1997, no. 6, (1997), 286–299.
- [12] P. Deift and X. Zhou, A steepest descent method for oscillatory Riemann–Hilbert problems, asymptotics for the mKDV equation, *Ann. Math.* 137 (1993), 295–368.
- [13] P. Deift and X. Zhou, Asymptotics for the Painlevé II equation. *Comm. Pure Appl. Math.* 48 (1995), 277–337.
- [14] T.M. Dunster, Uniform asymptotic expansions for the reverse generalized Bessel polynomials, and related functions, *SIAM J. Math. Anal.* 32 (2001), 987–1013.
- [15] A.S. Fokas, A.R. Its, and A.V. Kitaev, The isomonodromy approach to matrix models in 2D quantum gravity, *Commun. Math. Phys.* 147 (1992), 395–430.

- [16] A.A. Gonchar and E.A. Rakhmanov, Equilibrium distributions and the rate of rational approximation of analytic functions, *Mat. Sbornik* 134 (1987), 306–352. English translation in *Math. USSR-Sbornik* 62 (1989), 305–348.
- [17] E. Grosswald, Bessel polynomials, *Lecture Notes Math.* 698, Springer, New York, 1978.
- [18] S. Kamvissis, K.T-R McLaughlin, and P. Miller, Semiclassical soliton ensembles for the focusing nonlinear Schrödinger equation, manuscript.
- [19] T. Kriecherbauer and K.T-R McLaughlin, Strong asymptotics of polynomials orthogonal with respect to Freud weights. *Internat. Math. Res. Notices* 1999, no. 6, (1999), 299–333.
- [20] A. Martínez-Finkelshtein, P. Martínez-González, and R. Orive, On asymptotic zero distribution of Laguerre and generalized Bessel polynomials with varying parameters, *J. Comput. Appl. Math.* 127 (2001), 255–266.
- [21] F.W.J. Olver, The asymptotic expansion of Bessel functions of large order, *Philos. Trans. Roy. Soc. London Ser. A* 247 (1954), 328–368.
- [22] F.W.J. Olver, *Asymptotics and special functions*, Academic Press, New York, 1974. Reprinted by AK Peters, Wellesley, 1997.
- [23] O. Perron, *Die Lehre von den Kettenbrüchen*, 3rd ed. Teubner, Stuttgart, 1957.
- [24] I.E. Pritsker and R.S. Varga, The Szegő curve, zero distribution and weighted approximation, *Trans. Amer. Math. Soc.* 349 (1997), 4085–4105.
- [25] E.B. Saff and R.S. Varga, On the zeros and poles of Padé approximants to e^z , *Numer. Math.* 25 (1975), 1–14.
- [26] E.B. Saff and R.S. Varga, On the zeros and poles of Padé approximants to e^z . III, *Numer. Math.* 30 (1978), 241–266.
- [27] H. Stahl, Orthogonal polynomials with complex valued weight function, I and II, *Constr. Approx.* 2 (1986), 225–240, 241–251.
- [28] K. Strebel, *Quadratic differentials*, Springer-Verlag, Berlin, 1984.
- [29] G. Szegő, Über eine Eigenschaft der Exponentialreihe, *Sitzungsber. Berl. Math. Ges.* 23 (1924), 50–64.
- [30] G. Szegő, *Orthogonal polynomials*, AMS Colloquium Publications 23, Amer. Math. Soc., Providence RI, 1939.
- [31] R. Wong and J.-M. Zhang, Asymptotic expansions of the generalized Bessel polynomials, *J. Comput. Appl. Math.* 85 (1997), 87–112.

A.B.J. Kuijlaars

arno@wis.kuleuven.ac.be

Department of Mathematics, Katholieke Universiteit Leuven, Celestijnenlaan 200 B, 3001 Leuven, Belgium

K.T-R McLaughlin

mcl@math.unc.edu

Department of Mathematics, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA

and

mcl@math.arizona.edu

Department of Mathematics, University of Arizona, Tucson, AZ 85721, USA